

Printed Version of

Indian Internet Journal of Forensic Medicine & Toxicology

A Peer Reviewed Quarterly Journal

(Abstracted in Indian Science Abstracts, Indian Citation Index,
Google Scholar and Covered by Index Copernicus)



Founder Editor : **Prof. T.D. Dogra**

Editor : **Dr. Mukesh Yadav**



Print ISSN: 0973-1970
Online ISSN: 0974-4487

IndianJournals.com
A product of Diva Enterprises Pvt. Ltd.

Indian Internet Journal of Forensic Medicine and Toxicology

Vol. 19, No. 2, April-June 2021

Editorial Board	Contents	Page No.
Founder Prof. Dogra TD	<i>Original Articles</i>	
Editor Dr. Mukesh Yadav	<i>Role of Plasma Cholinesterase level in Assessment of Severity and Outcome in Organophosphorus Poisoning</i> <i>Mustaria Pankaj Kumar Shushil, Nirmal Arya, Mukesh Kumar Bansal and Ajay Agarwal</i>	23-26
Assistant Editors Dr. Lalwani S Dr. Srivastav PC	<i>Profile of Medico-legal Cases in Casualty of a Tertiary Care Hospital in Delhi</i> <i>Jitendra Pratap Singh, Mukesh Kumar Bansal and Mukesh Yadav</i>	27-31
Members Prof. Javed Usmani Prof. PC Dikshit Prof. SK Verma Prof. Dalbir Singh Prof. Atul Murari Prof. RK Gorea Prof. Ashok Srivastav Prof. Nagesh G Rao Prof. NK Agarwal Prof. OP Murty Prof. Gautum Biswas	<i>Epidemiology and Outcome of Death Due to Burn Injuries –A Prospective Study</i> <i>Surendra Singh, Archana Kaul and Rajesh Kumar Rai</i>	32-36
	<i>Case Reports</i>	
	<i>Pregnancy as a Result of Rape Presumed to Constitute a Grave Injury to the Mental Health of the Pregnant Woman: MTP Allowed: Chhattisgarh High Court</i> <i>Mukesh Yadav and Mukesh Kumar Bansal</i>	37-39
	<i>Failure to take X-Ray leading to Misdiagnosis: A Case of Medical Negligence: NCDRC</i> <i>Mukesh Yadav and Mukesh Kumar Bansal</i>	40-45
	<i>Rape Victim Suffering from Mental Retardation and Psychosis: MTP Allowed Without Consent: Kerala High Court</i> <i>Mukesh Yadav</i>	46-49

ICFMT, its editor and publisher disclaim responsibility and liability for any statement of fact and opinion, originality of contents and of any copyright violations by the author.

Printed & Published by: Diva Enterprises Pvt. Ltd. on behalf of Indian Congress of Forensic Medicine and Toxicology **Printed at** Spectrum, 208 A/14A, Savitri Nagar, New Delhi 110 017, **Published at** Diva Enterprises Pvt. Ltd., B-9, A-Block, L.S.C., Naraina Vihar, New Delhi 110028, India, **Editor** Dr. Mukesh Yadav

Original Article

Role of Plasma Cholinesterase level in Assessment of Severity and Outcome in Organophosphorus Poisoning

Mustaria Pankaj Kumar Shushil¹, Nirmal Arya², Mukesh Kumar Bansal^{3*} and Ajay Agarwal⁴

¹Assistant Professor, Department of Forensic Medicine, MLB Medical College, Jhansi, Uttar Pradesh, India

²Demonstrator, Department of Pharmacology, Govt. Allopathic Medical College, Banda, Uttar Pradesh, India

³Assistant Professor, Department of Forensic Medicine, Govt. Allopathic Medical College, Banda, Uttar Pradesh, India

⁴Professor & Head, Department of Forensic Medicine, S.N. Medical College, Agra, Uttar Pradesh, India

*Corresponding author email id: drmukesh1987@gmail.com

Received: 23-03-2021; Accepted: 14-05-2021

ABSTRACT

Introduction: Organophosphorus poisoning is associated with high rate of mortality if not diagnosed and treated in time. Therefore aim of present study is to evaluate the significance of cholinesterase activity in assessing severity and to predict the outcome of Organophosphorus poisoning. **Material and Methods:** Present study was conducted in the department of Forensic Medicine, S.N. Medical College & Hospital, Agra during the period of 18 months (February 2011 to July 2012). Total 160 patients were enrolled after applying inclusion and exclusion criteria. Peradeniya Organophosphorus Poisoning Scale was applied to all study subjects. **Result and Observation:** Majority of patients were married (70%) and male (64%). About 42% of cases belong to the age group 21-30 years. About 1/3rd of the cases had low socioeconomic status and about 1/3rd cases were farmers by occupation. On applying Peradeniya Organophosphorus Poisoning (POP) Scale, about one third of patients had tachypnoea, followed by impaired consciousness (29%). Serum/ plasma cholinesterase level showed higher mortality in patients having suppressed cholinesterase activity (< 4000 IU/L). **Conclusion:** Organophosphorus poisoning diagnosed at early stages of poisoning determining pseudocholinesterase activity form a reliable diagnostic test for management and can serve as lives saving tool.

Keywords: Organophosphorus poisoning, Plasma cholinesterase, Pseudocholinesterase, Tachypnoea

INTRODUCTION

Organophosphorus compound (OPCs) poisoning is a global problem and it is associated with high mortality rate if left undiagnosed and untreated in time ^[1]. Organophosphorus compounds are extensively used in developing countries like India as insecticides to increase production of agricultural commodities. Therefore developing countries like India report alarming rates of

toxicity and deaths due to OPCs poisoning ^[2]. There are approximately 3 million pesticide poisoning cases worldwide annually and more than 2,20,000 deaths. Soni *et al.* also detected Organophosphorus compound as most common poison on chemical analysis of viscera ^[3].

Monocrotophos, dimethoate, parathion, malathion, diazinon, etc. are most commonly used OPCs. These compounds can be absorbed by various routes like

inhalation, ingestion, dermal absorption etc. Diagnosis of OPCs poisoning is made on the basis of history, clinical examinations and biochemical investigations. Treatment of OPCs is done with antidotes particularly atropine, oximes, benzodiazepines and selective determination^[4].

OPCs are not only powerful inhibitors of cholinesterase but also act directly on cholinergic nerve endings^[2]. These compounds inhibit the enzyme acetylcholinesterase leading to accumulation of acetylcholine, which binds to muscarinic and nicotinic receptors throughout nervous system. Signs and symptoms of poisoning are due to persistent acetylcholine hyperstimulation at muscarinic and nicotinic receptors sites.^[4] The major forms of cholinesterase exist in vertebrates which can hydrolyze acetylcholine are plasma cholinesterase (Pseudo- or Butyryl or Serum Cholinesterase) and RBC cholinesterase (True, Specific Cholinesterase). Both these types of enzymes are inhibited by OPCs poisoning.

Therefore the aim of present study is to evaluate the significance of cholinesterase activity in assessing severity and to predict the outcome of Organophosphorus poisoning.

MATERIAL AND METHODS

Present study was conducted in department of Forensic Medicine in collaboration with the emergency department of S.N. Medical College & Hospital, Agra during the period of 18 months (February 2011 to July 2012). There were 240 patients of OP poisoning who were admitted through OPD and emergency during the study period. Among them, 160 patients were enrolled after applying inclusion and exclusion criteria.

Inclusion Criteria

A known case of organophosphorus poisoning, patient with clinical signs and symptoms suggestive of organophosphorus poisoning, patient showing evidence of organophosphorus poisoning on evaluation.

Exclusion Criteria

Patient brought dead or unknown poisoning, patient on Ventilator, patient with multiple poisoning with drugs such

as opioids, diazepam, barbiturates etc. and patients who received partial treatment outside and referred to our hospital were excluded.

A detailed history, clinical examination and relevant biochemical investigations were performed. A thorough clinical examination was carried out with particular reference to vital parameters, pupil size, assessment of central nervous system, respiratory system, cardiovascular system. This examination took place during initial resuscitation and treatment of the patient. Peradeniya Organophosphorus Poisoning Scale was applied to all study subjects and severity of OP poisoning was graded as mild, moderate and severe.

In all study subjects, 3 ml, of blood was collected on admission before administration of atropine and plasma cholinesterase was estimated. Plasma cholinesterase was estimated by colorimetric method using a kit provided by "Raichem of USA". The instrument used was RA – 50. 3 ml of plain blood was drawn and 5 u/ml of blood was centrifuged at 3000 rpm for 5 minutes. The serum of the patient was taken and added to the tube containing 1.55 ml of the reagent. Required information is collected on open ended semi-structural scheduled and compiled analysis with the help of MS Excel software.

RESULT AND OBSERVATIONS

Out of 160 patients, 84 (54%) patients had PChe levels more than 50% within normal range. Only 11 patients (7%) had severe poisoning with PChe levels less than 10% (Table 1).

Table 1: Distribution of cases according to Pseudo Cholinesterase (PChe) Levels

Plasma cholinesterase level (% of normal)	Number	Percentage
Normal	86	54
20 – 50	56	35
10 – 20	7	4
<10	11	7
Total	160	100

Table 2: Correlation between Plasma cholinesterase Levels and Outcome

Plasma cholinesterase levels (% of normal)	Survived (%)	Expired (%)
Normal	84 (98)	2 (2)
20 – 50	38 (68)	18 (32)
10 – 20	5(71)	2 (29)
<10	7(63)	4 (37)
Total	134 (84)	26 (16)

Table 2 shows correlation between Plasma cholinesterase Levels and outcome. Majority of patients 86 (54%) had subclinical poisoning with their PChe levels being >50%, out of which only 2 (2%) patients expired. Patients with PChe levels <50% had more mortality compared to patients with PChe levels >50%.

Table 3: Severity of Organophosphorus Poisoning based on Cholinesterase Levels (SCE) on Presentation (Day 1)

Severity	Range of SCE (IU/L)	Number of case	Percentage
Severe	<1000	11	7
Moderate	1000-2000	7	4
Mild	2001-3000	24	16
Mild	3001-4000	18	11
Mild	4001-5000	14	8
Mild	>5000	86	54
Total		160	100

(As per modified Proud foot Classification)

*The reference lab value for serum /plasma cholinesterase in the study: 5100 to 11700 IU. Table 3 shows severity of Organophosphorus poisoning based on cholinesterase levels (SCE) on day one Presentation. 7% cases showed SCE level < 1000 IU/L and classified as severe cases while 4% cases showed SCE level between 1000-2000 IU/L and classified as moderate. Cases showing SCE level > 2000 IU/L are classified as mild cases and majority (89%) cases belong to this category.

Table 4: Relation between Initial Level of Serum Cholinesterase and Mortality

Severity	Range of SCE (IU/L)	No. of cases (%)	Mortality (%)
Severe	< 1000	11 (7)	4 (36)
Moderate	1000-2000	7 (4)	2 (29)
Mild	2001-3000	24 (16)	9 (37)
Mild	3001-4000	18 (11)	7 (39)
Mild	4001-5000	14 (8)	2 (14)
Mild	> 5000	86 (54)	2 (2)
Total		160 (100)	26 (16)

(As per modified Proud foot Classification)

Table 4 shows relation between initial level of serum cholinesterase and mortality. Serum /plasma cholinesterase level shows higher mortality in patients having suppressed cholinesterase activity (< 4000 IU/L). The least rates of mortality were in the >5000 IU/L (2%).

DISCUSSION

In the present study, the majority of patients were in the age group of 21-30 years (42%), followed by the age group of 31-40 years (22%). Male to female ratio in present study is 1.75:1 and married to unmarried ratio in present study is 2.33:1. 90 (56%) patients belonged to rural areas and 70 (44%) belonged to urban areas. 58 (37%) patients were from upper lower-socio economic groups, followed by 44(27%) from lower class, in contrast to only 10 (6%) from upper class. On the basis of occupation, 61 (38%) patients were farmers, followed by 35 (22%) housewives and 26 (16%) students.

In the present study, vomiting and salivation was the commonest symptom (94%) followed by sweating (92%) and lacrimation (64%). The characteristic odor was present in 94% of patients. The other common clinical signs were tachypnoea (84%), altered consciousness (72%) and miosis (54%). Results of our study are in accordance with the studies done by Goel *et al.* [5], Kumar *et al.* [6] and Ramani *et al.* [7].

In the present study, 65% patients had mild grade, 32% had moderate grade and 3% patients severe grade of poisoning. The individual components of Peradeniya organophosphorus poisoning (POP) scale namely miosis, fasciculation, respiratory rate, bradycardia and level of consciousness were compared and results are similar with the study done by Goel *et al.* [5].

Plasma cholinesterase (PChe) levels were classified according to Proudfoot classification into subclinical (normal), mild, moderate and severe poisoning. In present study, 11(7%) patients had PChe level less than 10% of normal at admission indicating severe poisoning and 4 (37%) of patients expired. 7 (4%) patients had PChe level between 10-20% indicating moderate poisoning and 2 (29%) of patients were expired. 56 (35%) patients had PChe level between 20-50% indicating mild poisoning and 18 (32%) of patients were expired (Dreisbach's grading). Patients with PChe levels <50% of normal had more mortality compared with PChe levels >50% of normal. Hence it can be inferred that low plasma cholinesterase activity can be taken as a good diagnostic test for OP poisoning.

Recent studies by Ram *et al.* [8] showed that PChe activity below 10% of normal were associated with poor prognosis. Our findings are consistent with the studies done by various authors (Namba *et al.* [9]; Wadia *et al.* [10]; Ramani *et al.* [7] and Cunha *et al.* [11] who showed that plasma cholinesterase activity estimation is a reliable diagnostic test in OPCs poisoning and considered it a valid marker between severity of poisoning and prognosis of patients.

CONCLUSION

Organophosphorus poisoning is one of the most common modes of suicidal deaths in our country. By estimation of pseudo cholinesterase activity, early diagnosis of OP poisoning can be done. Farmers and other people involved in spraying insecticides should be educated regarding prophylactic measures to prevent accidental organophosphorus poisoning.

REFERENCES

- [1] Aardema H, Meertens JHJM, Lightenberg JJM *et al.* Organophosphorus pesticide poisoning: Cases and developments. *The Netherlands Journal of Medicine.* 2018;66(4):149-153.
- [2] Dua A. Serum cholinesterase levels in organophosphorus poisoning-does it Predict Mortality. *Journal of Associate Physicians India.* 2001;49:168-169.
- [3] Soni A, Bansal MK, Garg P, Yadav M. Socio-demographic Profile of Poisoning Cases- an Autopsy based Study. *Indian Internet Journal of Forensic Medicine & Toxicology.* 2019;17(1):5-11.
- [4] Aggarwal NK, Aggarwal BBL. Trends of poisoning in Delhi. *Journal of Indian Academy of Forensic Medicine.* 1998; 20(2): 32-35.
- [5] Goel A, Joseph S, Dutta TK. Organophosphate poisoning: Predicting the need for ventilator support. *Journal of Associate Physicians India.* 1998;46:786-790.
- [6] Kumar APN, Murthy GL, Rajshekhar L. Clinical profile of Organophosphate and carbamate poisoning-NIMS experience. *Journal of the Association of Physicians of India.* 2001;49:169-170.
- [7] Ramani *et al.* Serum cholinesterase levels as a indicator of prognosis in organophosphorus poisoning. *Journal of the Association of Physicians of India.* 1988; 36:23-24.
- [8] Ram JS, Kumar SS, Jayarajan A, Kuppuswamy G. Continuous infusion of high doses of atropine in the management of Organophosphorus compound poisoning. *Journal of Association Physicians India.* 1991;39:190-193.
- [9] Namba T, Nolte CT, Jackrel J, Grob D. Poisoning due to organophosphate insecticides. Acute and chronic manifestations. *The American Journal of Medicine.* 1971; 50:475-492.
- [10] Wadia RS, Sadagopan C, Amin RS, Sardesai HV. Neurological manifestations of Organophosphorus insecticide poisoning. *Journal of Neurol Neurosurg Psychiatry.* 1974;37:841-847.
- [11] Cunha AF, Felipe ISA *et al.* Neuroreflex control of cardiovascular function is impaired after acute poisoning with Chlorpyrifos, an Organophosphorus insecticide: possible short and long term clinical complications. *Toxicology.* 2018;398-99:13-22.

How to cite this article: Shushil MPK, Arya N, Bansal MK and Agarwal A. Role of Plasma Cholinesterase level in Assessment of Severity and Outcome in Organophosphorus Poisoning. *Indian Internet Journal of Forensic Medicine & Toxicology* 2021; 19(2): 23-26.